

FRAMED AND ORIENTED LINKS OF CODIMENSION 2

JIANHUA WANG

ABSTRACT. Sanderson [12] gave an isomorphism $\theta : \pi_m(\bigvee_{i=1}^r S_i^2) \longrightarrow \pi_m(\bigvee_{i=1}^{r+1} \mathbb{C}P_i^\infty)$. In this paper we construct for any subset $\sigma \subset \{1, 2, \dots, r\}$ an isomorphism θ_σ from $\pi_m(\bigvee_{i=1}^r S_i^2)$ to $\pi_m(\bigvee_{i=1}^{r+1} \mathbb{C}P_i^\infty)$. The inclusion $S^2 \vee S^2 \hookrightarrow \mathbb{C}P^\infty \vee \mathbb{C}P^\infty$ induces a homomorphism $f : \pi_m(S^2 \vee S^2) \longrightarrow \pi_m(\mathbb{C}P^\infty \vee \mathbb{C}P^\infty)$. We also compute f by evaluating f on each factor in the Hilton splitting of $\pi_m(S^2 \vee S^2)$, the results in [12] concerning the case $m = 4$ are generalized.

Keywords: Framed and oriented links, Seifert surface, Hilton splitting.

2000 Mathematics Subject Classification: 55Q20, 57Q45

1. INTRODUCTION

A link $M_1 \sqcup M_2 \sqcup \dots \sqcup M_r \subset \mathbb{R}^m$ is an ordered disjoint union of closed smooth submanifolds, $m \geq 3$. If these submanifolds are oriented then we call it an oriented link. A framing of a k -codimensional submanifold in $\mathbb{R}^m$ is a trivialization of its normal vector bundle, or equivalently, an ordered set of k linearly independent normal vector fields. If every component of the link has a given framing then we call it a framed link. We will assume that the codimensions of the components are 2, when nothing else is stated. The bordism groups of framed and oriented links with r components of codimension 2 in $\mathbb{R}^m$ are denoted by $FL_{m,r}^2$ and $L_{m,r}^2$ respectively. The facts

$$FL_{m,r}^2 \cong \pi_m(\bigvee_{i=1}^r S_i^2), \quad L_{m,r}^2 \cong \pi_m(\bigvee_{i=1}^r \mathbb{C}P_i^\infty)$$

are well known by Pontryagin-Thom construction.

The main purpose of this paper is to discuss the relationship between $FL_{m,r}^2$ and $L_{m,r+1}^2$ and to compute the homomorphism

$$f : \pi_m(S^2 \vee S^2) \longrightarrow \pi_m(\mathbb{C}P^\infty \vee \mathbb{C}P^\infty) \cong \pi_m(S^2).$$

Geometrically, f is given by forgetting the framing of $[M_1 \sqcup M_2]_{fr} \in FL_{m,2}^2$ and keeping the orientation determined by the framing, so we may call f a forgetful homomorphism.

Main results and the organization of this paper: In §2 we give a formula of reframing, suggested by Koschorke, and discuss briefly the role of framing in the Hilton splitting. Let $\sigma \subset \{1, 2, \dots, r\}$ be any subset. We construct in §3 an isomorphism $\theta_\sigma : FL_{m,r}^2 \longrightarrow L_{m,r+1}^2$. We recover Sanderson's isomorphism θ by taking $\sigma = \{1, 2, \dots, r\}$. The forgetful homomorphism f is computed in §4 by using the inverse of θ_σ and by

choosing $\sigma = \phi$, in particular the following result of [12] is generalized: in case $m = 4$ it holds $f \circ \gamma_* \neq 0$ for $\gamma = [\iota_1, \iota_2]$, $[\iota_1, [\iota_1, \iota_2]]$ and $[\iota_2, [\iota_1, \iota_2]]$, $f \circ \gamma_* = 0$ for $\gamma = \iota_1$ and ι_2 .

We work in the category of smooth manifolds.

I'm grateful to my supervisor Prof. U. Koschorke for some ideas and stimulating discussions, and to Prof. U. Kaiser for many helps and useful suggestions. Thanks also to Prof. M. Heusener for nice talks. As the English version of this paper is finished in Bar-Ilan University I am also thankful to Prof. T. Nowik for being warmly hosted.

2. FRAMING

Let $M^{m-2} \subset \mathbb{R}^m$ be a closed submanifold with framing $\mathcal{F} = (v_1, v_2)$, and let

$$s : M \longrightarrow S^1 \simeq SO(2)$$

be a continuous map. For $x \in M$ we can represent $s(x) \in S^1$ by an orthonormal matrix

$$\begin{pmatrix} a_{11}(x) & a_{12}(x) \\ a_{21}(x) & a_{22}(x) \end{pmatrix}.$$

Define v'_1 and v'_2 by

$$\begin{aligned} v'_1(x) &= a_{11}(x)v_1(x) + a_{12}(x)v_2(x), \\ v'_2(x) &= a_{21}(x)v_1(x) + a_{22}(x)v_2(x). \end{aligned}$$

(v'_1, v'_2) is a new framing of M and is denoted by $s\mathcal{F}$. Up to homotopy we may assume s is differential. Let $-1 \in S^1$ be a regular value of s and consider $Z = s^{-1}(-1) \subset M$. Let $Z \times [-1, 1] \subset M$ be a small tubular neighbourhood of Z such that the positive direction of $[-1, 1]$ is in agreement with the usual orientation of S^1 . Up to homotopy $s\mathcal{F}$ is in fact the 2π -rotation of $\mathcal{F}$ in this neighbourhood, namely outside this neighbourhood it is the same as (v_1, v_2) and inside it

$$v'_1(z, t) = v_1(z, t) \cos(t+1)\pi + v_2(z, t) \sin(t+1)\pi, \quad (1)$$

$$v'_2(z, t) = -v_1(z, t) \sin(t+1)\pi + v_2(z, t) \cos(t+1)\pi, \quad (2)$$

where $z \in Z$ and $t \in [-1, 1]$.

Let v_3 be the normal vector field of $Z \subset M$, determined by the orientation of S^1 , provide $Z \subset \mathbb{R}^m$ with the framing (v_1, v_2, v_3) . We define

$$[Z, \mathcal{F}, s] = [Z, (v_1, v_2, v_3)] \in \pi_m(S^3).$$

Let $\eta : S^3 \longrightarrow S^2$ be the Hopf map, Koschorke observed that $[Z, \mathcal{F}, s]$ is the only obstruction to homotope $\mathcal{F}$ to $s\mathcal{F}$ and conjectured that $\eta_*[Z, \mathcal{F}, s]$ and the difference

$$[M, s\mathcal{F}] - [M, \mathcal{F}] \in \pi_m(S^2)$$

should be related by some formula.

It's well known that M bounds a Seifert surface F . If the framing of M induced by F is homotopic to the given framing $\mathcal{F}$ of M , then we say M is S -framed. In this case we have $[M, \mathcal{F}] = 0$, and Turaev [15] proved $[M, s\mathcal{F}] = \eta_*[Z, \mathcal{F}, s]$. Note that if M is oriented then the S -framing compatible with the orientation is unique up to homotopy.

Proposition 2.1. *Let $M \subset \mathbb{R}^m$ be a closed submanifold with framing $\mathcal{F} = (v_1, v_2)$ and $s : M \rightarrow S^1$ be a map. Let $u\mathcal{F} = (-v_1, v_2)$ and M^{sh} be a small shift of M along v_1 provided with the framing $s\mathcal{F} = (v'_1, v'_2)$. Then it holds*

$$[(M \sqcup M^{sh}), (u\mathcal{F} \sqcup s\mathcal{F})] = \eta_*[Z, \mathcal{F}, s].$$

Proof. Writing $(M \sqcup M^{sh})$ we mean it is considered as submanifold rather than a link of two components. Without loss of generality we may assume that $\mathcal{F} = (v_1, v_2)$ is smooth and orthogonal. Let $\hat{W} \cong M \times [0, 1]$ be the trace of a shift from M to M^{sh} along v_1 . Cut out a small ε -neighbourhood U of $Z \times \{\frac{1}{2}\} \subset \hat{W}$, and define $W = \hat{W} \setminus U$, where $Z = s^{-1}(-1)$. See Fig.1.




FIGURE 1.

Provide $W \subset \mathbb{R}^m \times \{0\} \subset \mathbb{R}^{m+1}$ with the framing $\mathcal{G} = (e_{m+1}, v_2)$, where e_{m+1} is the last vector in the usual base of $\mathbb{R}^{m+1}$. Let $Z \times [-\varepsilon, \varepsilon] \subset M$ be a ε -neighbourhood of Z . Up to homotopy we may assume s maps $M \setminus Z \times [-\varepsilon, \varepsilon]$ to the base point $1 \in S^1$. We get now a well defined map $\hat{s} : W \rightarrow S^1$, given by

$$\hat{s}(x, t) = \begin{cases} s(x) & : t \geq \frac{1}{2}, \\ 1 & : t \leq \frac{1}{2}. \end{cases}$$

So we obtain a new framing $\hat{s}\mathcal{G}$ of $W \subset \mathbb{R}^{m+1}$. In addition, it holds $\partial W = M \sqcup M^{sh} \sqcup \partial U$, where $\partial U \cong Z \times S^1$ is the boundary of U . We construct now a diffeotopy of $\mathbb{R}^{m+1}$ which deforms $(W, \hat{s}\mathcal{G})$ to a framed bordism.

Let $\nu(M^{sh})$ be the normal vector bundle of $M^{sh} \subset \mathbb{R}^{m+1}$, framed by (v_1, e_{m+1}, v_2) . A homotopy of $\nu(M^{sh})$

$$F_1 : \nu(M^{sh}) \times [0, 1] \rightarrow \nu(M^{sh})$$

is given by rotating e_{m+1} to v_1 , v_1 to $-e_{m+1}$ and meanwhile keeping v_2 fixed. Define

$$F'_1 : M^{sh} \times [0, 1] \longrightarrow \mathbb{R}^{m+1}$$

by $F'_1(x, t) = (x, -t)$. Let U_1 be a δ -neighbourhood of $M^{sh} \subset \mathbb{R}^{m+1}$ with $\delta \ll \epsilon$. From F_1, F'_1 we get an isotopy $H_1 : U_1 \times [0, 1] \longrightarrow \mathbb{R}^{m+1}$, given by

$$\begin{aligned} & H_1(x + r_1 v_1(x) + r_2 e_{m+1} + r_3 v_2(x), t) \\ &= F'_1(x, t) + r_1 F_1(v_1(x), t) + r_2 F_1(e_{m+1}, t) + r_3 F_1(v_2(x), t), \end{aligned}$$

where $x \in M^{sh}$ and $x + r_1 v_1(x) + r_2 e_{m+1} + r_3 v_2(x) \in U_1$. It holds clearly

$$H_1(M^{sh}, 1) = M^{sh} \times \{-1\} \subset \mathbb{R}^m \times \{-1\}.$$

Since (e_{m+1}, v_2) is deformed to (v_1, v_2) and $\hat{s}|_{M^{sh}} = s$ (s is defined on M^{sh} by identifying M^{sh} with M in the natural way), $\hat{s}\mathcal{G}|_{M^{sh}}$ is homotoped to $s\mathcal{F}$.




FIGURE 2.

Let U_0 be a δ -neighbourhood of $M \subset \mathbb{R}^{m+1}$. Similarly we have an isotopy

$$H_0 : U_0 \times [0, 1] \longrightarrow \mathbb{R}^{m+1}$$

which deforms M to $M \times \{-1\} \subset \mathbb{R}^m \times \{-1\}$ and (e_{m+1}, v_2) to $(-v_1, v_2) = u\mathcal{F}$. We have used the homotopy $F_0 : \nu(M) \times [0, 1] \longrightarrow \nu(M)$ which rotates e_{m+1} to $-v_1$ and keeps v_2 fixed.

Let $\nu(\partial U)$ be the normal vector bundle of $\partial U \subset \mathbb{R}^{m+1}$, framed by (u_1, e_{m+1}, v_2) , where u_1 is the normal vector field of $\partial U \subset W$ pointing inwards. Let

$$F_2 : \nu(\partial U) \times [0, 1] \longrightarrow \nu(\partial U)$$

be the homotopy given by rotating e_{m+1} to u_1 and keeping v_2 fixed; and define

$$F'_2 : \partial U \times [0, 1] \longrightarrow \mathbb{R}^{m+1}$$

by $F'_2(x, t) = (x, t)$. From F_2 and F'_2 we obtain an isotopy $H_2 : U_2 \times [0, 1] \longrightarrow \mathbb{R}^{m+1}$, where U_2 is a δ -neighbourhood of $\partial U \subset \mathbb{R}^{m+1}$. H_2 isotopes ∂U to $\partial U \times \{1\} \subset \mathbb{R}^m \times \{1\}$ and homotopes (e_{m+1}, v_2) to (u_1, v_2) . So $\hat{s}\mathcal{G}|_{\partial U}$ is homotoped to $\hat{s}(u_1, v_2)$. It is not difficult to see that $\hat{s}(u_1, v_2)$ is homotopic to the 2π -rotation of the S -framing (u_1, v_2) of $\partial U \subset \mathbb{R}^m \times \{1\}$, and therefore

$$(\partial U, \hat{s}(u_1, v_2)) = (Z \times S^1, \hat{s}(u_1, v_2))$$

is just the fibre-wise embedding of the framed circle $S^1 \subset \mathbb{R}^3$ representing $\eta : S^3 \longrightarrow S^2$. It follows $[\partial U, \hat{s}(u_1, v_2)] = \eta_*[Z, \mathcal{F}, s]$.

Now $U_\delta = U_0 \cup U_1 \cup U_2$ is a δ -neighbourhood of $\partial W \subset \mathbb{R}^{m+1}$. The isotopies H_0 , H_1 and H_2 together define an isotopy $H : U_\delta \times [0, 1] \longrightarrow \mathbb{R}^{m+1}$. According to a well known theorem in Differential Topology H determine a diffeotopy $\tilde{H}$ of $\mathbb{R}^{m+1}$ which deforms $(W, \hat{s}\mathcal{G})$ to a framed bordism from $[(M \sqcup M^{sh}), (u\mathcal{F} \sqcup s\mathcal{F})]$ to $\eta_*[Z, \mathcal{F}, s]$, see Fig.2. $\square$

Corollary 2.2. (i) [Turaev, 1985]: *It holds $[M, s\mathcal{F}] = \eta_*[Z, \mathcal{F}, s]$, if $\mathcal{F}$ is the S -framing of the submanifold M ;*

(ii) $E[M, s\mathcal{F}] - E[M, \mathcal{F}] = E\eta_*[Z, \mathcal{F}, s]$, where E is the suspension homomorphism.

Proof. (i) Let F be a Seifert surface of M giving the S -framing $\mathcal{F}$. Because M^{sh} is a small shift of M along $\mathcal{F}$, we have $M^{sh} \cap F = \emptyset$. Using F and $M^{sh} \times [0, 1]$ we obtain easily a framed bordism from $[(M \sqcup M^{sh}), (u\mathcal{F} \sqcup s\mathcal{F})]$ to $[M, s\mathcal{F}]$. The assertion follows now from Proposition 2.1. See also Turaev [15].

(ii) We have clearly

$$\begin{aligned} E[(M \sqcup M^{sh}), (u\mathcal{F} \sqcup s\mathcal{F})] &= E[M, u\mathcal{F}] + E[M, s\mathcal{F}] \\ &= -E[M, \mathcal{F}] + E[M, s\mathcal{F}]. \end{aligned}$$

The statement follows from Proposition 2.1. $\square$

In general it is a subtle problem to measure the difference $[M, s\mathcal{F}] - [M, \mathcal{F}]$. Hilton-Hopf invariant up to order 3 are involved. Consider the framed link $(M, u\mathcal{F}) \sqcup (M^{sh}, s\mathcal{F})$ representing an element α in

$$\pi_m(S_1^2 \vee S_2^2) \cong \oplus_\gamma \pi_m(S^{q(\gamma)+1}),$$

where γ runs through a system Γ of basic Whitehead products in $\iota_1 < \iota_2$, and $q(\gamma)$ is the height of $\gamma \in \Gamma$, see Hilton [2]. So we have the splitting $\alpha = \oplus_\gamma \alpha_\gamma$ with $\alpha_\gamma \in \pi_m(S^{q(\gamma)+1})$. Let $\alpha_1, \alpha_2, \alpha_3$ be the Hilton coefficients of α corresponding to $[\iota_1, \iota_2]$, $[\iota_1, [\iota_1, \iota_2]]$ and

$[\iota_2, [\iota_1, \iota_2]]$ respectively. If we map the wedge $S_1^2 \vee S_2^2$ canonically to the sphere S^2 , then ι_1 and ι_2 are both identified with the identity ι of S^2 , and α is mapped to the element

$$[(M \sqcup M^{sh}), (u\mathcal{F} \sqcup s\mathcal{F})] \in \pi_m(S^2).$$

The basic Whitehead products in $\iota_1 < \iota_2$ are mapped to Whitehead products in ι . Because the Whitehead products in ι with weight > 3 are zero homotopic, we have

$$\begin{aligned} & [(M \sqcup M^{sh}), (u\mathcal{F} \sqcup s\mathcal{F})] \\ = & [M, u\mathcal{F}] + [M^{sh}, s\mathcal{F}] + [\iota, \iota]_* \alpha_1 + [\iota, [\iota, \iota]]_* (\alpha_2 + \alpha_3). \end{aligned}$$

Let α'_1 be the first nontrivial Hilton-Hopf invariant of $[M, \mathcal{F}]$, namely the one corresponding to $[\iota_1, \iota_2]$, then it holds (see Hilton [2])

$$[M, u\mathcal{F}] = -[M, \mathcal{F}] + [\iota, \iota]_* \alpha'_1.$$

By this and Proposition 2.1 we get

Corollary 2.3. *It holds*

$$[M, s\mathcal{F}] - [M, \mathcal{F}] = \eta_*[Z, \mathcal{F}, s] - [\iota, \iota]_* (\alpha_1 + \alpha'_1) - [\iota, [\iota, \iota]]_* (\alpha_2 + \alpha_3).$$

Note that $3[\iota, [\iota, \iota]] = 0$ by Jacobi-identity.

Consider now a framed link $(M_1, \mathcal{F}_1) \sqcup (M_2, \mathcal{F}_2) \subset \mathbb{R}^m$ of codimensions $k_1, k_2 \geq 2$. The bordism group $FL_m^{k_1, k_2}$ of such links is isomorphic to $\pi_m(S^{k_1} \vee S^{k_2})$. We try to understand the role of the framings $\mathcal{F}_1, \mathcal{F}_2$ in the Hilton splitting of $(M_1, \mathcal{F}_1) \sqcup (M_2, \mathcal{F}_2)$.

Let $i = 1, 2$. A Seifert surface of M_i is a compact oriented submanifold $F_i \subset \mathbb{R}^m$ with boundary M_i . If M_i has a framed Seifert surface F_i such that the framing of M_i as the boundary of F_i is homotopic to the original framing $\mathcal{F}_i$, then we say $\mathcal{F}_i$ is an S -framing of M_i and M_i is S -framed. We call F_i a suitably framed Seifert surface. Note that two S -framings must not be homotopic.

Proposition 2.4. *Let $(M_1, \mathcal{F}_1) \sqcup (M_2, \mathcal{F}_2) \subset \mathbb{R}^m$ be an S -framed link representing $\alpha \in \pi_m(S^{k_1} \vee S^{k_2})$, and let F_1, F_2 be the corresponding suitably framed Seifert surfaces.*

(i) Up to involution it holds $[M_1 \natural F_2] = [F_1 \natural M_2]$. An involution is an isomorphism u of the target group with $u \circ u = id$.

(ii) Let $\alpha = \oplus_\gamma \alpha_\gamma$ be the Hilton splitting of α . Up to involution $\alpha_{[\iota_1, \iota_2]}$ is given by $[M_1 \natural F_2]$, all other Hilton coefficients α_γ are zero.

Proof. The desired framed bordism in the first part is given by $F_1 \natural F_2$.

Let $Z = M_1 \natural F_2$ and let $U \cong Z \times D^{k_1}$ be an open tubular neighbourhood of $Z \subset F_2$. Frame ∂U so that $F_2 \setminus U$ gives rise to a framed bordism between M_2 and ∂U . Because $F_2 \setminus U$ is disjoint from M_1 , $M_1 \sqcup M_2$ is framed bordant to $M_1 \sqcup \partial U$, see Fig.3.




FIGURE 3.

If U is small enough, then it holds $F_1 \cap \bar{U} \cong Z \times [0, 1]$ with $Z = Z \times \{0\}$. Let v be the normal vector field of $Z \subset F_1 \cap \bar{U}$ pointing inwards. It's easy to see that $Z' = Z \times \{1\} = F_1 \cap \partial U$ is just a small shift of Z along v . Let U' be a small tubular neighbourhood of $Z' \subset F_1$ such that $U' \cap \partial U = Z'$. Frame $\partial U'$ so that $F_1 \setminus U'$ gives rise to a framed bordism between M_1 and $\partial U'$. Because $F_1 \setminus U'$ is disjoint from ∂U it follows $[M_1 \sqcup \partial U] = [\partial U' \sqcup \partial U]$, see Fig.3 again.

In addition, $\partial U' \sqcup \partial U = Z \times S^{k_2-1} \sqcup Z \times S^{k_1-1}$ is just the fibre-wise embedding of the standard framed Hopf link $S^{k_2-1} \sqcup S^{k_1-1} \subset \mathbb{R}^{k_1+k_2-1}$ (at least up to involution of the framings) into a small tubular neighbourhood of the framed intersection $Z \subset \mathbb{R}^m$. It follows

$$[M_1 \sqcup M_2] = [\iota_1, \iota_2]_*[Z]$$

at least up to involution. Since the Hilton splitting of $[\iota_1, \iota_2]_*[Z]$ has the form

$$0 + 0 + [Z] + 0 + \cdots,$$

the assertion follows. $\square$

Corollary 2.5. *Let γ be a basic Whitehead product in $\iota_1 < \iota_2$ of weight ≥ 3 , and let $M_1 \sqcup M_2 \subset \mathbb{R}^{q(\gamma)+1}$ be a framed link representing γ . At least one component of this link is not S -framed.*

Proof. If this is not the case then according to the above result we have $H_\gamma[M_1 \sqcup M_2] = 0$, a contradiction to the fact $H_\gamma[M_1 \sqcup M_2] = \pm 1 \in \pi_{q(\gamma)+1}(S^{q(\gamma)+1}) \cong Z$, where H_γ is the Hilton homomorphism corresponding to the basic Whitehead product γ . $\square$

3. ISOMORPHISMS BETWEEN $FL_{m,r}^2$ AND $L_{m,r+1}^2$

Sanderson [12] gave an isomorphism $\theta : FL_{m,r}^2 \longrightarrow L_{m,r+1}^2$. We construct here for each subset $\sigma \subset \{1, 2, \dots, r\}$ such an isomorphism θ_σ .

Let $L = (M_1, \mathcal{F}_1) \sqcup \cdots \sqcup (M_r, \mathcal{F}_r) \subset \mathbb{R}^m$ be a framed link of codimension 2. For $i \in \sigma$ define $\hat{M}_i = -M_i^{sh}$, where $-M_i^{sh}$ is the negative oriented M_i^{sh} . Note that a

framed submanifold in $\mathbb{R}^m$ is canonically oriented. For $i \notin \sigma$ consider the framed intersection $Z_i = M_i \cap \tilde{F}_i$, where $\tilde{F}_i$ is a Seifert surface of M_i^{sh} . Let U_i be a small tubular neighbourhood of $Z_i \subset \tilde{F}_i$. If we orient ∂U_i as the boundary of U_i then $\tilde{F}_i \setminus U_i$ is a Seifert surface of $(-M_i^{sh}) \sqcup (-\partial U_i)$. For $i \notin \sigma$ we define $\hat{M}_i = \partial U_i$ and $M_{r+1} = \sqcup_{i=1}^r \hat{M}_i$. θ_σ is given by the assignment

$$\theta_\sigma[L] = [M_1 \sqcup \cdots \sqcup M_r \sqcup M_{r+1}]_{or}.$$

Lemma 3.1. $\theta_\sigma : FL_{m,r}^2 \longrightarrow L_{m,r+1}^2$ is a well defined homomorphism.

Proof. To prove this let $W_1 \sqcup \cdots \sqcup W_r \subset \mathbb{R}^m \times [0, 1]$ be a framed bordism between

$$L = \sqcup_{i=1}^r (M_i, \mathcal{F}_i), \quad L' = \sqcup_{i=1}^r (M'_i, \mathcal{F}'_i).$$

Take $\tilde{W}_i = W_i^{sh}$ and let $\tilde{F}_i, \tilde{F}'_i$ be Seifert surfaces of M_i^{sh} and $M'_i{}^{sh}$. We define for $i \notin \sigma$

$$\hat{W}_i = \tilde{W}_i \cup \tilde{F}_i \times \{0\} \cup \tilde{F}'_i \times \{1\}$$

and orient $\hat{W}_i$ so that its orientation is in agreement with $\tilde{W}_i$. Let $F_{\hat{W}_i}$ be a Seifert surface of $\hat{W}_i$, consider $\hat{Z}_i = W_i \cap F_{\hat{W}_i}$ with boundary $\partial \hat{Z}_i = Z_i \times \{0\} \sqcup Z'_i \times \{1\}$. Cut out a small tubular neighbourhood $\hat{U}_i$ of $\hat{Z}_i \subset F_{\hat{W}_i}$ and orient $\partial \hat{U}_i$ so that $\partial \hat{U}_i$ is an oriented bordism from ∂U_i to $\partial U'_i$. Define

$$W_{r+1} = (\sqcup_{i \notin \sigma} \partial \hat{U}_i) \sqcup (-\sqcup_{i \in \sigma} W_i^{sh}).$$

$W_1 \sqcup \cdots \sqcup W_r \sqcup W_{r+1}$ is clearly an oriented bordism between $\theta_\sigma(L)$ and $\theta_\sigma(L')$. It is clear that θ_σ respects the addition. $\square$

To prove that θ_σ is an isomorphism we construct now a homomorphism $\zeta_\sigma : L_{m,r+1}^2 \longrightarrow FL_{m,r}^2$ and show it's in fact the inverse of θ_σ .

Let $\bar{\sigma} = \sigma \cup \{r+1\}$, and let $L = M_1 \sqcup \cdots \sqcup M_r \sqcup M_{r+1} \subset \mathbb{R}^m$ be an oriented link of codimension 2. Take a Seifert surface $F_{\bar{\sigma}}$ of $M_{\bar{\sigma}} = \sqcup_{i \in \bar{\sigma}} M_i$. Denote by u_1 the normal vector field of $M_{\bar{\sigma}} \subset F_{\bar{\sigma}}$ pointing outwards and by u_2 the normal vector field of $F_{\bar{\sigma}}$ determined by the orientation. For $i \in \sigma$ take $\mathcal{F}_i = (u_1, u_2)|_{M_i}$ as a framing of M_i . If $i \notin \sigma$ consider the intersection $Z_i = M_i \cap F_{\bar{\sigma}}$ and let $Z_i \times [-1, 1] \subset M_i$ be a small tubular neighbourhood of $Z_i \subset M_i$ such that the positive direction of $[-1, 1]$ is in agreement with $u_2|_{Z_i}$. We assume here that M_i intersects $F_{\bar{\sigma}}$ perpendicularly. Let (v_1^S, v_2^S) be the S -framing of M_i determined by the orientation. Define $\mathcal{F}_i = (v_1, v_2)$ to be the 2π -rotation of (v_1^S, v_2^S) in $Z_i \times [-1, 1]$, see (1) and (2) in §2. By doing this we have provided M_i with the framing $\mathcal{F}_i$ for $1 \leq i \leq r$. Now $\zeta_\sigma : L_{m,r+1}^2 \longrightarrow FL_{m,r}^2$ is defined by the following assignment

$$\zeta_\sigma[M_1 \sqcup \cdots \sqcup M_r \sqcup M_{r+1}]_{or} = [(M_1, \mathcal{F}_1) \sqcup \cdots \sqcup (M_r, \mathcal{F}_r)].$$

Lemma 3.2. ζ_σ is a well defined homomorphism.

Proof. Let $W_1 \sqcup \cdots \sqcup W_r \sqcup W_{r+1} \subset \mathbb{R}^m \times [0, 1]$ be an oriented bordism between

$$L = M_1 \sqcup \cdots \sqcup M_r \sqcup M_{r+1}, \quad L' = M'_1 \sqcup \cdots \sqcup M'_r \sqcup M'_{r+1}$$

and let F'_σ be a Seifert surface of $M'_\sigma = \sqcup_{i \in \sigma} M'_i$. Consider

$$W = W_{\bar{\sigma}} \cup F_{\bar{\sigma}} \times \{0\} \cup F'_\sigma \times \{1\},$$

where $W_{\bar{\sigma}}$ is the disjoint union of the W_i 's with $i \in \bar{\sigma}$. We orient W so that its orientation coincides with the one of $W_{\bar{\sigma}}$. Smooth W and let F_W be an oriented Seifert surface of W . F_W induces canonically a framing $\mathcal{G}_i$ on W_i for each $i \in \sigma$. If $i \notin \sigma$ consider

$$\hat{W}_i = W_i \cup F_i \times \{0\} \cup F'_i \times \{1\},$$

where F_i, F'_i are Seifert surfaces of M_i and M'_i , and again we orient $\hat{W}_i$ so that its orientation is in agreement with the one of W_i . Let $\mathcal{G}_i^S$ be the restriction of the S -framing of $\hat{W}_i$ on W_i . Consider now the intersection $\hat{Z}_i = W_i \cap F_W$ with boundary

$$\partial \hat{Z}_i = Z_i \times \{0\} \sqcup Z'_i \times \{1\},$$

and denote by $\hat{Z}_i \times [-1, 1] \subset W_i$ a small tubular neighbourhood such that the positive direction of $[-1, 1]$ coincides with the normal vector field of $F_W \subset \mathbb{R}^{m+1}$ determined by the orientation. For $i \notin \sigma$ let $\mathcal{G}_i$ be the framing of W_i given by the 2π -rotation of $\mathcal{G}_i^S$ in $\hat{Z}_i \times [-1, 1]$. Clearly

$$(W_1, \mathcal{G}_1) \sqcup \cdots \sqcup (W_r, \mathcal{G}_r)$$

is a framed bordism from $\zeta_\sigma[L]$ to $\zeta_\sigma[L']$, and ζ_σ evidently respects the addition. $\square$

Theorem 3.3. $\theta_\sigma : FL_{m,r}^2 \longrightarrow L_{m,r+1}^2$ is an isomorphism, in fact its inverse is ζ_σ .

Proof. (1) $\zeta_\sigma \circ \theta_\sigma = id$. Consider

$$\begin{aligned} \zeta_\sigma \circ \theta_\sigma[(M_1, \mathcal{F}_1) \sqcup \cdots \sqcup (M_r, \mathcal{F}_r)] &= \zeta_\sigma[M_1 \sqcup \cdots \sqcup M_r \sqcup M_{r+1}]_{or} \\ &= [(M_1, \mathcal{F}'_1) \sqcup \cdots \sqcup (M_r, \mathcal{F}'_r)]. \end{aligned}$$

For M_{r+1} and the framings $\mathcal{F}'_i$ see the definitions of θ_σ and ζ_σ . Denote by $M_i \times [0, \varepsilon]$ the trace of a small ε -shift from M_i to M_i^{sh} . Orient

$$F_{\bar{\sigma}} = (\sqcup_{i \notin \sigma} U_i) \sqcup (\sqcup_{i \in \sigma} M_i \times [0, \varepsilon])$$

so that $F_{\bar{\sigma}}$ is a Seifert surface of $M_{\bar{\sigma}}$. According to the definition of ζ_σ we see easily that up to homotopy it holds $\mathcal{F}'_i = \mathcal{F}_i$ for $i \in \sigma$. In addition, for $i \notin \sigma$ the intersection $M_i \cap F_{\bar{\sigma}} = M_i \cap U_i$ is exactly the submanifold $Z_i = M_i \cap \tilde{F}_i$, because U_i is a small tubular neighbourhood of $Z_i \subset \tilde{F}_i$, see the definition of θ_σ . This implies that $\mathcal{F}_i$ and $\mathcal{F}'_i$ are homotopic, since both $\mathcal{F}_i$ and $\mathcal{F}'_i$ are essentially the 2π -rotation of the S -framing of M_i in a small tubular neighbourhood of $Z_i \subset M_i$. It follows $\zeta_\sigma \circ \theta_\sigma = id$.

(2) $\theta_\sigma \circ \zeta_\sigma = id$. Consider

$$\begin{aligned} \theta_\sigma \circ \zeta_\sigma [M_1 \sqcup \cdots \sqcup M_r \sqcup M_{r+1}]_{or} &= \theta_\sigma [(M_1, \mathcal{F}_1) \sqcup \cdots \sqcup (M_r, \mathcal{F}_r)] \\ &= [M_1 \sqcup \cdots \sqcup M_r \sqcup M'_{r+1}]_{or}. \end{aligned}$$

For M'_{r+1} and the framings see the definitions of θ_σ and ζ_σ . Let $F_{\bar{\sigma}}$ be an oriented Seifert surface of $M_{\bar{\sigma}}$. Cut out a small tubular neighbourhood V_σ of $M_\sigma \subset F_{\bar{\sigma}}$ to get F'_σ with

$$\partial F'_\sigma = \partial(F_{\bar{\sigma}} \setminus V_\sigma) = M_{r+1} \sqcup (\sqcup_{i \in \sigma} M_i^{sh}).$$

For $i \notin \sigma$ let $\tilde{F}_i$ be an oriented Seifert surface of M_i^{sh} . According to the definitions of θ_σ and ζ_σ we may assume

$$Z_i = M_i \cap \tilde{F}_i = M_i \cap F'_\sigma$$

for $i \notin \sigma$, because $\partial \tilde{F}_i = M_i^{sh}$ is a shift of M_i along the framing $\mathcal{F}_i$ ($\mathcal{F}_i$ is given by the 2π -rotation of the S -framing in a small tubular neighbourhood of $M_i \cap F'_\sigma \subset M_i$), and because $M_i \cap \tilde{F}_i$ is just where the 2π -rotation of the S -framing takes place. In addition, we may assume M_i intersects $\tilde{F}_i$ perpendicularly. This implies $U_i \subset F'_\sigma$ after a small isotopy of F'_σ fixing boundary, where $U_i \subset \tilde{F}_i$ is a small tubular neighbourhood of $Z_i = M_i \cap \tilde{F}_i$. Define

$$\hat{F}_\sigma = F'_\sigma \setminus (\cup_{i \notin \sigma} U_i)$$

which is a Seifert surface of

$$M_{r+1} \sqcup (\sqcup_{i \in \sigma} M_i^{sh}) \sqcup (\sqcup_{i \notin \sigma} -\partial U_i) = M_{r+1} \sqcup (-M'_{r+1}).$$

Because $\hat{F}_\sigma$ is disjoint from all M_i , $1 \leq i \leq r$, there is an embedding $W_{r+1} \subset \mathbb{R}^m \times [0, 1]$ of $\hat{F}_\sigma$ such that W_{r+1} is an oriented bordism from M_{r+1} to M'_{r+1} and such that W_{r+1} is disjoint from $M_i \times [0, 1] \subset \mathbb{R}^m \times [0, 1]$ for all $1 \leq i \leq r$. So we obtain an oriented bordism

$$M_1 \times [0, 1] \sqcup \cdots \sqcup M_r \times [0, 1] \sqcup W_{r+1} \subset \mathbb{R}^m \times [0, 1]$$

from $M_1 \sqcup \cdots \sqcup M_r \sqcup M_{r+1}$ to $M_1 \sqcup \cdots \sqcup M_r \sqcup M'_{r+1}$. It follows $\theta_\sigma \circ \zeta_\sigma = id$. $\square$

Example 3.4. If $\sigma, \sigma' \subset \{1, \dots, r\}$ are different then $\theta_\sigma \neq \theta_{\sigma'}$ in general. Without loss of generality we assume $1 \in \sigma$ and $1 \notin \sigma'$. Consider the framed Hopf link

$$L = S_1^1 \sqcup S_2^1 \sqcup \phi \sqcup \cdots \sqcup \phi \subset \mathbb{R}^3.$$

Let $L_\sigma, L_{\sigma'}$ be the oriented links representing $\theta_\sigma[L]$ and $\theta_{\sigma'}[L]$ respectively. It is not difficult to verify that the linking number between the second and the last components of L_σ is ± 1 , and the linking number between the second and the last components of $L_{\sigma'}$ is 0. This shows $\theta_\sigma \neq \theta_{\sigma'}$. In the case $\sigma = \{1, \dots, r\}$ it holds $\theta_\sigma = \theta$, where θ is the isomorphism given by Sanderson [12]. Some computations may be simplified by suitably choosing θ_σ or ζ_σ . In next section we will choose $\zeta_\sigma = \zeta_\phi$, namely $\sigma = \phi$.

4. COMPUTATION OF THE FORGETFUL HOMOMORPHISM

Let $f : FL_{m,2}^2 \longrightarrow L_{m,2}^2$ be the forgetful homomorphism given by forgetting the framings. To compute f we compute the composition

$$f' = \zeta_\sigma \circ f : FL_{m,2}^2 \longrightarrow L_{m,2}^2 \xrightarrow{\cong} FL_{m,1}^2$$

where and throughout this section $\sigma = \phi$.

Let $M^{m-2} \subset \mathbb{R}^m$ be a framed or an oriented submanifold. Denote by M^S the same submanifold but provided with the S -framing. Let $M_1 \sqcup M_2 \subset \mathbb{R}^m$ be a framed link representing an element in $FL_{m,2}^2$. According to the definition of ζ_σ and from Corollary 2.2 we see

$$f'[M_1 \sqcup M_2]_{fr} = \eta_*[M_1^S \natural F_2]_{fr}$$

at least up to involution, where $\eta : S^3 \longrightarrow S^2$ is the Hopf map and F_2 is a Seifert surface of M_2 . So we need only to compute $M_1^S \natural F_2$.

It is easily seen that $f' \circ \gamma_* = 0$ for $\gamma = \iota_1$ and $\gamma = \iota_2$. So let γ be a basic Whitehead product in $\iota_1 < \iota_2$ of weight ≥ 2 , and $\eta_\gamma : S^{q(\gamma)+1} \longrightarrow S^3$ be the map determined by $M_1^S(\gamma) \natural F_2(\gamma)$, where $M_1(\gamma) \sqcup M_2(\gamma)$ is a framed link representing γ and $F_2(\gamma)$ is a framed Seifert surface of $M_2(\gamma)$. Given $[Z] \in \pi_m(S^{q(\gamma)+1})$, by fibre-wise embedding it follows easily that $f' \circ \gamma_*[Z] = \eta_\gamma[Z]$ which is represented by $Z \times (M_1^S(\gamma) \natural F_2(\gamma))$. Therefore we only need to compute $M_1^S(\gamma) \natural F_2(\gamma)$.

For $\gamma = [\iota_1, \iota_2]$ we see easily that $M_1^S(\gamma) \natural F_2(\gamma)$ is a framed point, this means η_γ is the identity up to sign in this case. Let $\gamma_w = [\iota_2, [\iota_2 \cdots [\iota_2, [\iota_1, \iota_2]] \cdots]]$ be a basic Whitehead product of weight $w \geq 3$, clearly $q(\gamma_w) = w$. The following framed link in $\mathbb{R}^{w+1}$ represents γ_w

$$\begin{aligned} M_1(w) &= S_w^1 \times S_{w-1}^1 \times \cdots \times S_3^1 \times S_2^1, \\ M_2(w) &= S_w^1 \times S_{w-1}^1 \times \cdots \times S_3^1 \times S_1^1 \sqcup \\ &\quad S_w^1 \times S_{w-1}^1 \times \cdots \times S^2 \sqcup \\ &\quad \dots \sqcup \\ &\quad S_w^1 \times S^{w-2} \sqcup \\ &\quad S^{w-1} \\ &= N_{2,1} \sqcup \cdots \sqcup N_{2,w-1}, \end{aligned}$$

where $S_2^1 \sqcup S_1^1$, $S_i^1 \sqcup S^{i-1}$ are S -framed Hopf links, $3 \leq i \leq w$, and all products are given by fibre-wise embeddings. $M_1(w)$ is clearly S -framed, but $M_2(w)$ not, according to Corollary 2.5. In addition, we can assume $M_1(w) \subset \mathbb{R}^w \times \{0\} \subset \mathbb{R}^{w+1}$.

Consider now the oriented submanifolds of $M_1(w)$

$$Z_i = S_w^1 \times \cdots \times S_{i+1}^1 \times \{pt\} \times S_{i-1}^1 \cdots \times S_3^1 \times S_2^1,$$

$2 \leq i \leq w$ and $\{pt\}$ is a set of a single point. $\cup_{i=2}^w Z_i \subset M_1(w)$ is a transversally immersed submanifold. Using the classical trick in Fig.4 we can successively dissolve the multi-points to get an embedded submanifold $Z(w) \subset M_1(w)$ of codimension 1. Let v_1, v_2 be the normal vector fields of $Z(w) \subset M_1(w)$ and of $M_1(w) \subset \mathbb{R}^w \times \{0\}$ respectively. Define $\mathcal{F}_w = (v_1, v_2|_{Z(w)})$.



FIGURE 4.




FIGURE 5.

For the proof of Proposition 4.1 we need to consider the following situation. Let

$$N^{m-2} = N_1 \sqcup N_2 \subset \mathbb{R}^m$$

be an oriented closed submanifold and let F_1, F_2 be Seifert surfaces of N_1 and N_2 respectively. We construct now a Seifert surface F of N from F_1 and F_2 . Cut out

tubular neighborhoods of $N_1 \subset F_1$ and $N_2 \subset F_2$ to get F_1^* and F_2^* , and cut out a tubular neighbourhood $U(C)$ of

$$C = F_1^* \cap F_2^* \subset F_1^* \cup F_2^*.$$

Then we can use the trick in Fig.5 to sew $F_1^* \setminus U(C)$ and $F_2^* \setminus U(C)$ together to get F^* with $\partial F^* = N_1^{sh} \sqcup N_2^{sh}$, where N_1^{sh} is essentially a small shift of N_1 along the framing given by the 2π -rotation of the S -framing of N_1 in a tubular neighbourhood of $N_1 \cap F_2 \subset N_1$, similarly N_2^{sh} . Let T_1, T_2 be the traces of these shifts. $F^* \cup T_1 \cup T_2$ is oriented, we can make it smooth and get a Seifert surface F of N . Note that, shown in Fig.5 is the locus near $F_1 \cap N_2$, for the locus $N_1 \cap F_2$ it is completely similar; away from $F_1 \cap N_2$ and $N_1 \cap F_2$ the method shown in Fig.4 applies. For details see [16], p.14–17.

Proposition 4.1. (a) Let $\gamma_w, M_1(w), M_2(w), Z(w)$ and $\mathcal{F}_w$ be as above, and assume $w \geq 3$. There exists a Seifert surface $F_2(w)$ of $M_2(w)$ such that the following holds

(i) $M_1(w) \cap F_2(w) = Z(w),$

(ii) $[M_1^S(w) \cap F_2(w)] = \pm E[Z(w), \mathcal{F}_w].$

(b) Up to sign the framed submanifold $M_1^S(w) \cap F_2(w) \subset \mathbb{R}^{w+1}$ represents the map

$$\eta_\gamma = (E\eta) \circ (E^2\eta) \circ \cdots \circ (E^{w-3}\eta) \circ (E^{w-2}\eta) : S^{w+1} \longrightarrow S^3.$$

Proof. (a) The case $w = 3$ is easy. Assume inductively that the assertion is true for γ_{w-1} , and let $F_2(w-1)$ be a Seifert surface of $M_2(w-1)$ with the desired property.

By fibre-wise embedding we get a Seifert surface $S_w^1 \times F_2(w-1)$ of $N_{2,1} \sqcup \cdots \sqcup N_{2,w-2} \cdot N_{2,w-1} = S^{w-1}$ bounds a ball D^w . From D^w and $S_w^1 \times F_2(w-1)$ we obtain a Seifert surface $F_2(w)$ of $M_2(w)$. To compute $M_1(w) \cap F_2(w)$ we look at

$$\begin{aligned} Z' &= M_1(w) \cap S_w^1 \times F_2(w-1) \\ &= S_w^1 \times (M_1(w-1) \cap F_2(w-1)) \\ &= S_w^1 \times Z(w-1), \\ Z'' &= M_1(w) \cap D^w \\ &= \{pt\} \times S_{w-1}^1 \times \cdots \times S_3^1 \times S_2^1 \\ &= Z_w. \end{aligned}$$

Considered in $M_1(w)$ we have the transversal intersection

$$Q = Z' \cap Z'' = \{pt\} \times Z(w-1)$$

of codimension 2. Because Q is disjoint from the boundaries of $S_w^1 \times F_2(w-1)$ and D^w we see that in the construction of $F_2(w)$ we have just dissolved Q as Fig.4. This means $M_1(w) \cap F_2(w) = Z(w)$. Part (i) follows. Because $M_1(w)$ is S -framed, at least up to sign $\mathcal{F}_w$ is the framing of $M_1^S(w) \cap F_2(w) = Z(w)$, part (ii) follows.

(b) Up to involution we have

$$[M_1^S(w) \cap F_2(w)] = [F_1(w) \cap M_2^S(w)],$$

see Proposition 2.4. Consider

$$\begin{aligned} Z'(w) &= F_1(w) \cap M_2(w) \\ &= F_1(w) \cap N_{2,1} \\ &= S_w^1 \times S_{w-1}^1 \times \cdots \times S_3^1 \times D_2 \cap \\ &\quad S_w^1 \times S_{w-1}^1 \times \cdots \times S_3^1 \times S_1^1, \\ &= S_w^1 \times S_{w-1}^1 \times \cdots \times S_3^1 \times \{pt\}, \end{aligned}$$

where D_2 is a disk with boundary S_2^1 . Let (v_1, v_2) be the S -framing of $M_2(w)$ and v_3 be the normal vector field of $F_1(w) \subset \mathbb{R}^{w+1}$. Define $\mathcal{F}'_w = (v_1, v_2, v_3)|_{Z'(w)}$. So we have

$$[Z'(w), \mathcal{F}'_w] = [F_1(w) \cap M_2^S(w)],$$

In addition, $(v_1, v_2)|_{N_{2,1}}$ is given by the 2π -rotations of the S -framing (u_1, u_2) of $N_{2,1}$ in the tubular neighborhoods of all

$$S_w^1 \times \cdots \times S_{i+1}^1 \times \{pt\} \times S_{i-1}^1 \times \cdots \times S_3^1 \times S_1^1 \subset N_{2,1},$$

$3 \leq i \leq w$ and $\{pt\}$ denotes a set of a single point. We will get $(Z'(w), \mathcal{F}'_w)$ when we take a regular value in S^3 of the map

$$(E\eta) \circ (E^2\eta) \circ \cdots \circ (E^{w-3}\eta) \circ (E^{w-2}\eta) : S^w \longrightarrow S^3$$

and perform the Pontryagin-Thom construction. The statement follows. $\square$

Proposition 4.2. *Let $\gamma = [\iota_2, [\iota_2, \cdots [\iota_2, [\iota_1, [\iota_1, \cdots [\iota_1, \iota_2] \cdots]]] \cdots]$ be a basic Whitehead product in $\iota_1 < \iota_2$ of weight $w \geq 3$, and let $M_1 \sqcup M_2 \subset \mathbb{R}^{w+1}$ be a framed link representing γ . The following map*

$$\eta_\gamma = (E\eta) \circ (E^2\eta) \circ \cdots \circ (E^{w-3}\eta) \circ (E^{w-2}\eta) : S^{w+1} \longrightarrow S^3$$

is represented by $M_1^S \cap F_2$ up to sign, where F_2 is a Seifert surface of M_2 .

Proof. Assume ι_i appears w_i -times in γ , $i = 1, 2$. The following framed link $M_1 \sqcup M_2$ in $\mathbb{R}^{w+1}$ representing γ

$$\begin{aligned} M_1 &= S_w^1 \times S_{w-1}^1 \times \cdots \times S_{w_1+2}^1 \times S_{w_1+1}^1 \cdots \times S_3^1 \times S_2^1 \sqcup \\ &\quad S_w^1 \times S_{w-1}^1 \times \cdots \times S_{w_1+2}^1 \times S_{w_1+1}^1 \cdots \times S^2 \sqcup \\ &\quad \cdots \cdots \cdots \sqcup \\ &\quad S_w^1 \times S_{w-1}^1 \times \cdots \times S_{w_1+2}^1 \times S^{w_1} \\ &= N_{1,1} \sqcup \cdots \sqcup N_{1,w_1} \end{aligned}$$

$$\begin{aligned} M_2 &= S_w^1 \times S_{w-1}^1 \times \cdots \times S_{w_1+3}^1 \times S_{w_1+2}^1 \times \cdots \times S_3^1 \times S_1^1 \sqcup \\ &\quad S_w^1 \times S_{w-1}^1 \times \cdots \times S_{w_1+3}^1 \times S^{w_1+1} \sqcup \\ &\quad \cdots \cdots \cdots \sqcup \\ &\quad S_w^1 \times S^{w-2} \sqcup \\ &\quad S^{w-1} \\ &= N_{2,1} \sqcup \cdots \sqcup N_{2,w_2}. \end{aligned}$$

where $S_2^1 \sqcup S_1^1$, $S_3^1 \sqcup S^2, \dots$, $S_w^1 \sqcup S^{w-1}$ are usually framed Hopf links and all products are given by fibre-wise embeddings.

According to Proposition 4.1 we can assume

$$\begin{aligned} Z &= F_1 \natural M_2 = F_1 \natural N_{2,1} \\ &= S_w^1 \times S_{w-1}^1 \times \cdots \times S_{w_1+2}^1 \times Z(w_1 + 1), \end{aligned}$$

where $Z(w_1 + 1)$ is given by dissolving the multi-points of the following immersion iteratedly

$$\begin{aligned} &\cup_{i=1, i \neq 2}^{w_1+1} S_{w_1+1}^1 \times \cdots \times S_{i+1}^1 \times \{pt\} \times S_{i-1}^1 \times \cdots \times S_3^1 \times S_1^1 \\ &\subset S_{w_1+1}^1 \times \cdots \times S_3^1 \times S_1^1. \end{aligned}$$

Let $Z'(w_1 + 1) = S_{w_1+1}^1 \times \cdots \times S_3^1 \times \{pt\}$, framed by (v_1, v_2, v_3) as in the proof of Proposition 4.1, part (b). So we have a framed bordism $(W', \mathcal{G}')$ from $Z(w_1 + 1)$ to $Z'(w_1 + 1)$ (at least up to involution) with $\mathcal{G}' = (\bar{v}_1, \bar{v}_2, \bar{v}_3)$. Consider the fibre-wise embedding

$$W = S_w^1 \times \cdots \times S_{w_1+2}^1 \times W' \subset \mathbb{R}^{w+1} \times [0, 1].$$

We obtain a framing $\mathcal{G}$ of W by performing the 2π -rotations of $(\bar{v}_2, \bar{v}_3)$ in tubular neighborhoods of all

$$S_w^1 \times \cdots \times S_{i+1}^1 \times \{*\} \times S_{i-1}^1 \times \cdots \times S_{w_1+2}^1 \times W' \subset W,$$

$w_1 + 2 \leq i \leq w$. $(W, \mathcal{G})$ is a framed bordism from $F_1 \natural M_2^S$ to $Z'(w)$, according to the proof of Proposition 4.1, part (b), $Z'(w)$ represents

$$(E\eta) \circ (E^2\eta) \circ \cdots \circ (E^{w-3}\eta) \circ (E^{w-2}\eta) : S^{w+1} \longrightarrow S^3.$$

The assertion follows by Lemma 2.4. $\square$

Note that in case $m = 4$ it holds $f' \circ \gamma_* \neq 0$ for $\gamma = [\iota_1, \iota_2]$, $[\iota_1, [\iota_1, \iota_2]]$ and $[\iota_2, [\iota_1, \iota_2]]$, and $f' \circ \gamma_* = 0$ for $\gamma = \iota_1$ and ι_2 . Because ζ_ϕ is an isomorphism, the same holds if we replace f' by f . So we recover the corresponding results of Sanderson [12].

Let Γ be a system of basic Whitehead products in $\iota_1 < \iota_2$, and $\gamma = [\alpha, \beta] \in \Gamma$ be such that the weights of α and β are greater than 1. Take γ' to be one of α , β and γ , and let $M_1(\gamma') \sqcup M_2(\gamma') \subset \mathbb{R}^{q(\gamma')+1}$ be a framed link representing γ' . Denote by $\eta_{\gamma'} : S^{q(\gamma')+1} \longrightarrow S^3$ the map given by $[M_1^S(\gamma') \natural F_2(\gamma')]_{fr}$, where $F_2(\gamma')$ is a Seifert surface of $M_2(\gamma')$.

Proposition 4.3. *Under the above notations and up to sign we have*

$$\eta_\gamma = [\eta_\alpha, \eta_\beta] : S^{q(\alpha)+q(\beta)+1} = S^{q(\gamma)+1} \longrightarrow S^3.$$

Proof. Surely, we may take $M_1(\gamma) \sqcup M_2(\gamma) \subset \mathbb{R}^{q(\gamma)+1}$ to be the following

$$\begin{aligned} M_1(\gamma) &= S^{q(\alpha)} \times M_1(\beta) \sqcup S^{q(\beta)} \times M_1(\alpha), \\ M_2(\gamma) &= S^{q(\alpha)} \times M_2(\beta) \sqcup S^{q(\beta)} \times M_2(\alpha), \end{aligned}$$

where $S^{q(\alpha)} \sqcup S^{q(\beta)} \subset \mathbb{R}^{q(\gamma)+1}$ is the usually framed Hopf link and all products are given by fibre-wise embeddings. Let $F_i(\alpha)$, $F_i(\beta)$ be Seifert surfaces of $M_i(\alpha)$ and $M_i(\beta)$ respectively, $i = 1, 2$, then we get a Seifert surface

$$F_i(\gamma) = S^{q(\alpha)} \times F_i(\beta) \sqcup S^{q(\beta)} \times F_i(\alpha)$$

of $M_i(\gamma)$ by fibre-wise embeddings. This implies

$$\begin{aligned} M_1^S(\gamma) &= S^{q(\alpha)} \times M_1^S(\beta) \sqcup S^{q(\beta)} \times M_1^S(\alpha), \\ M_1^S(\gamma) \natural F_2(\gamma) &= S^{q(\alpha)} \times (M_1^S(\beta) \natural F_2(\beta)) \sqcup S^{q(\beta)} \times (M_1^S(\alpha) \natural F_2(\alpha)). \end{aligned}$$

Because by induction $M_1^S(\alpha) \natural F_2(\alpha)$, $M_1^S(\beta) \natural F_2(\beta)$ represent the maps η_α and η_β respectively, the assertion $\eta_\gamma = [\eta_\alpha, \eta_\beta]$ follows. $\square$

Combining Propositions 4.1, 4.2 and 4.3 we get a complete computation of f' and therefore the forgetful homomorphism f . Given $\alpha = \oplus_\gamma \alpha_\gamma \in \pi_m(S^2 \vee S^2)$, $f'(\alpha)$ is the sum of $f' \circ \gamma_*(\alpha_\gamma)$, and all the homomorphisms $f' \circ \gamma_*$ are computed above. To get the decomposition $\alpha = \oplus_\gamma \alpha_\gamma$ we may use the method in the author's paper [17]. We end this paper with a question which U. Kaiser mentioned to me. Given a framed link $(M_1, \mathcal{F}_1) \sqcup (M_2, \mathcal{F}_2) \subset \mathbb{R}^m$ of codimension 2, denote by $\mathcal{F}_i^S$ the S -framing of M_i ,

$i = 1, 2$. There is a map $s_i : M_i \longrightarrow S^1$ such that $\mathcal{F}_i \simeq s_i \mathcal{F}_i^S$. Let $q_i : S^1 \longrightarrow S^1$ be a map of degree q_i and define $s'_i = q_i \circ s_i$. The assignment

$$[(M_1, \mathcal{F}_1) \sqcup (M_2, \mathcal{F}_2)] \longmapsto [(M_1, s'_1 \mathcal{F}_1^S) \sqcup (M_2, s'_2 \mathcal{F}_2^S)]$$

determines a well defined homomorphism $(q_1, q_2)_* : FL_{m,2}^2 \longrightarrow FL_{m,2}^2$. The question is

How can one describe the homomorphism $(q_1, q_2)_*$ by using the Hilton splitting of $FL_{m,2}^2 \cong \pi_m(S^2 \vee S^2)$?

REFERENCES

- [1] J.S.Carter, S.Kamada, M.Saito, S.Satoh, A Theorem of Sanderson on Link Bordisms in Dimension 4, *Algebraic and Geometric Topology* **Vol 1**, 299-310 (2001)
- [2] P.J.Hilton, On the Homotopy Groups of the Union of Spheres, *J. London Math. Soc.* **30**, 154-172 (1955)
- [3] M.W.Hirsch, Differential Topology, Springer-Verlag, New York - Heidelberg - Berlin (1976)
- [4] U.Kaiser, Link Homotopy in $\mathbb{R}^3$ and S^3 , *Pacific J. of Math.* **vol 151, No.2**, 257-264 (1991)
- [5] U.Kaiser, Verschlingungsabbildungen im Euklidischen Raum, *Dissertation*, Uni. Siegen (1989)
- [6] M.A.Kervaire, An Interpretation of G.Whitehead's Generalization of H.Hopf's Invariant, *Ann. of Math.* **vol 69, No.2**, 335-365 (1959)
- [7] U.Koschorke, Higher Order Homotopy Invariants for Higher Dimensional Link Maps, *Lecture Notes in Mathematics 1172*, Springer-Verlag, 116-129 (1985)
- [8] U.Koschorke, Homotopy, Cocordance and Bordism of Link Maps, *Global Analysis in Modern Mathematics*, Publish or Perish, 283-299 (1993)
- [9] U.Koschorke, A Generalization of Milnor's μ -Invariants to Higher Dimensional Link Maps, *Topology* **vol 36, No.2**, 301-324 (1997)
- [10] U.Koschorke, P.A.Kirk, Generalized Seifert Surfaces and Linking Numbers, *Topology and Its Applications* **42, No.3**, 247-262 (1991)
- [11] L.S.Pontryagin, Smooth Manifolds and Their Applications in Homotopy Theory, *Amer. Math. Transaction* **vol (2) 2**, 1-114 (1959)
- [12] B.Sanderson, Bordism of Links in Codimension 2, *J. London Math. Soc.* **vol (2) 35**, 367-376 (1987)
- [13] B.Sanderson, Triple Links in Codimension 2, *Colloquia Mathematica Societatis Jano Bolyai* **vol 55**, PECS (Hungary), 457-471 (1989)
- [14] N.Sato, Cobordisms of Semi-boundary Links, *Topology and Its Applications* **vol 18**, 225-234 (1984)
- [15] V.Turaev, The Sato-Levine-Cochran Invariants Vanish for Higher Dimensional Links, preprint, Leningrad (1985)
- [16] J.Wang, Eine Geometrische Interpretation Gewisser Hilton-Koeffizienten, *Dissertation*, Uni. Siegen (1998)
- [17] J.Wang, The Geometry of The Hilton Splitting (to appear)
- [18] G.W.Whitehead, Elements of Homotopy Theory, Springer-Verlag, New York - Heidelberg - Berlin (1978)

Department of Mathematics
Bar-Ilan University
52900 Ramat Gan, Israel
e-mail: wangji@macs.biu.ac.il